

Some topics in complex and harmonic analysis, 4

Stephen William Semmes
Rice University
Houston, Texas

Let $\mathcal{S}(\mathbf{R}^n)$ denote the Schwartz class of rapidly decreasing smooth complex-valued functions on $\mathbf{R}^n$, and let $\mathcal{S}'(\mathbf{R}^n)$ denote the vector space of tempered distributions on $\mathbf{R}^n$, which are the continuous linear functionals on $\mathcal{S}(\mathbf{R}^n)$. Let $\phi \in \mathcal{S}(\mathbf{R}^n)$ and $\lambda \in \mathcal{S}'(\mathbf{R}^n)$ be given, and for each $x \in \mathbf{R}^n$ let ϕ_x be the function defined by

$$(1) \quad \phi_x(y) = \phi(x - y).$$

Thus $\phi_x \in \mathcal{S}(\mathbf{R}^n)$ for all $x \in \mathbf{R}^n$.

The convolution of λ and ϕ is the function on $\mathbf{R}^n$ defined by

$$(2) \quad (\lambda * \phi)(x) = \lambda(\phi_x).$$

One can check that this is in fact a continuous function on $\mathbf{R}^n$. More precisely, $x \mapsto \phi_x$ is continuous as a function defined on $\mathbf{R}^n$ with values in the Schwartz class, and hence $\lambda(\phi_x)$ is continuous as a complex-valued function on $\mathbf{R}^n$ because λ is a continuous linear functional on the Schwartz class. One can show further that $\lambda * \phi$ is a smooth function on $\mathbf{R}^n$.

For each pair of multi-indices α, β we have the semi-norm $\|\cdot\|_{\alpha, \beta}$ defined on the Schwartz class by

$$(3) \quad \|\psi\|_{\alpha, \beta} = \sup \left\{ \left| x^\alpha \frac{\partial^{d(\beta)}}{\partial x^\beta} \psi(x) \right| : x \in \mathbf{R}^n \right\},$$

where $d(\beta)$ denotes the degree of $\beta = (\beta_1, \dots, \beta_n)$, which is defined to be the sum $\beta_1 + \dots + \beta_n$. This family of seminorms determines the topology on the Schwartz class. In particular, the statement that λ is a continuous linear functional on the Schwartz class means that there is a nonnegative real number C and multi-indices $\alpha_1, \beta_1, \dots, \alpha_p, \beta_p$ such that

$$(4) \quad |\lambda(\psi)| \leq C \sum_{j=1}^p \|\psi\|_{\alpha_j, \beta_j}$$

for all functions ψ in the Schwartz class.

As a result, one can show that $\lambda * \phi$ is a function of polynomial growth on $\mathbf{R}^n$. To be more precise, this means that there is a nonnegative real number C_1 and a nonnegative integer l such that

$$(5) \quad |(\lambda * \phi)(x)| \leq C_1 (1 + |x|)^l$$

for all $x \in \mathbf{R}^n$. That is, $|\lambda(\phi_x)|$ is bounded by a constant times a finite sum of seminorms applied to ϕ_x , and one can check that these seminorms have polynomial growth in x . One can take this further and show that the derivatives of $\lambda * \phi$ also have polynomial growth.

It is perhaps better to say that $\lambda * \phi$ has at most polynomial growth. If $\lambda(\psi)$ is defined by integrating ψ times a function in the Schwartz class, for instance, then $\lambda * \phi$ is the same as a convolution of two functions in the Schwartz class, and is an element of the Schwartz class. For that matter, one can show that $\lambda * \phi$ lies in the Schwartz class if λ has compact support, or if λ is defined by integration by a measure with finite total mass.

As another interesting special case, suppose that $\lambda(\psi)$ is defined by integrating ψ times a fixed polynomial $p(x)$ on $\mathbf{R}^n$. The definition of the convolution $(\lambda * \phi)(x)$ is the same as the integral of $\phi(x - y)$ times $p(y)$ with respect to y , and this is equal to the integral of $\phi(y)$ times $p(x - y)$ with respect to y . One can expand $p(x - y)$ as a linear combination of products of monomials in x and y , and the monomials in x can be taken outside the integral. Thus $\lambda * \phi$ is a polynomial, with the coefficients given by moments of ϕ , which is to say integrals of ϕ times monomials. It is a simple matter to write down examples with nonvanishing coefficients of whatever degree one might like.

By definition, a polynomial is a linear combination of monomials x^α . A polynomial of degree at most l is a linear combination of monomials x^α where the multi-indices α have degree at most l . A homogeneous polynomial of degree l is a linear combination of monomials x^α where the multi-indices α have degree equal to l . This is equivalent to saying that the polynomial $p(x)$ is homogeneous of degree l in the sense that $p(tx) = t^l p(x)$ for all $x \in \mathbf{R}^n$ and all positive real numbers t , or arbitrary real numbers t for that matter. By differentiating $p(tx)$ in t and evaluating at $t = 1$ we get the well-known identity that $\sum_{j=1}^n x_j (\partial/\partial x_j) p(x)$ is equal to l times $p(x)$ when $p(x)$ is homogeneous of degree l .

A polynomial $p(x)$ is said to be harmonic if $\Delta p = 0$, where Δ denotes the usual Laplace operator, which is to say the sum of $\partial^2/\partial x_j^2$ over $j = 1, \dots, n$. If p is a polynomial of degree at most l , then p can be written in a canonical way as the sum of polynomials which are homogeneous of degrees ranging from 0 to l . For if p is given as a linear combination of monomials x^α where the multi-indices α have degree less than or equal to l , one can simply group the terms together of the same degree. If p is a harmonic polynomial, then the homogeneous components of p are also harmonic. For any polynomial p , the Laplacian applied to the homogeneous components of p of degree 0 or 1 is equal to 0, and the Laplacian applied to a homogeneous component of p of degree $m \geq 2$ is the same as the homogeneous component of the polynomial Δp of degree $m - 2$.

Let p be a harmonic polynomial on $\mathbf{R}^n$. We would like to check that the average of p over the unit sphere in $\mathbf{R}^n$ is equal to the value of p at the origin. It is enough to check this when p is a harmonic polynomial which is homogeneous of some degree l , since we can express any harmonic polynomial as a sum of homogeneous harmonic polynomials. When $l = 0$, our homogeneous harmonic

polynomial is simply a constant, and thus its average over the unit sphere is equal to its value at the origin. Now suppose that $l \geq 1$, in which case we would like to show that the average of p over the unit sphere is equal to 0.

By the identity mentioned earlier, if x is a unit vector in $\mathbf{R}^n$, then $p(x)$ is equal to a nonzero constant times the derivative of p at x in the direction of the unit normal to the unit sphere. The divergence theorem tells us that the integral of the unit normal derivative of p on the unit sphere is equal to 0, because the integral of the Laplacian of p on the unit ball is equal to 0, since p is harmonic. Therefore the integral of p on the unit sphere is equal to 0, as desired.

This shows that the average of a harmonic polynomial on the unit sphere is equal to the value of the polynomial at the origin. In general the average of a harmonic polynomial over any sphere is equal to the value of the harmonic polynomial at the center of the sphere. Using this one can verify that the convolution of a harmonic polynomial p with a radial function ϕ with suitable integrability properties, such as a radial function in the Schwartz class, is equal to the integral of ϕ over $\mathbf{R}^n$ times p . In other words, harmonic polynomials are eigenfunctions for suitable radial convolution operators.

References

- [1] W. Rudin, *Functional Analysis*, second edition, McGraw-Hill, 1991.
- [2] E. Stein and S. Shakarchi, *Fourier Analysis: An Introduction*, Princeton Lectures in Analysis **1**, Princeton University Press, 2003.
- [3] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Mathematical Series **32**, Princeton University Press, 1971.